

Spectral Distortions of the CMB

In this problem, you will investigate whether a proposed satellite mission, the Primordial Inflation Experiment (PIXIE, Kogut et al., 2011, 2016), will be able to detect spectral distortions of the Cosmic Microwave Background (CMB).

Over the past 30 years, the CMB has provided research opportunities for *precision cosmology*¹, the results of which have solidified our understanding of the history of our universe. The spatial anisotropies of the CMB have been at the forefront of both experiment and theory, and have driven the field forward. As per our understanding, these spatial anisotropies can reveal details about the size and physical properties of the primordial fluctuations in matter distribution leading to the rise of the structure that we see today. The interaction of matter with photons, on the other hand should lead to spectral anisotropies. Such anisotropies have been theoretically predicted, but have never been detected so far.

In 1992, the FIRAS instrument on NASA's Cosmic Background Explorer (COBE) (Fixsen et al., 1996; Mather et al., 1994; Fixsen, 2009) mission showed that the CMB spectrum is a near-perfect blackbody, isotropic to the level of one part in 10^5 , with temperature $T_{\text{CMB}} = (2.725 \pm 0.001)$ K.

Radiation is thermalized at very high redshifts (well before and up until the decoupling of radiation and matter) due to a number of radiative processes, such as Compton, inverse Compton, double Compton scattering and bremsstrahlung, each one dominating at different energy regimes (and thus redshifts).²

Physical processes such as reionization and structure formation, decaying or annihilating particles, dissipation of primordial density fluctuations, adiabatic cooling of matter and recombination, as well as cosmic strings, primordial black holes, small-scale magnetic fields, can lead to distortions to the blackbody shape of the spectrum of CMB (Chluba, 2014; Chluba and Jeong, 2014; Hill et al., 2015; Tashiro, 2014).

Spectral distortions of the CMB, if present, were not detectable at the precision level of the FIRAS instrument. PIXIE will provide a sensitivity improved by 76 times compared to its similar predecessor.

As the Universe expands, processes responsible for maintaining thermalization become less effective. This then allows spectral distortions to be developed. The distortions created between redshift $z \approx 10^6$ (when double Compton scattering decouples) and $z \approx 10^5$ (when Compton scattering no longer contributes to thermalization) are characterized by a chemical potential³ and are called " μ distortions" (Zeldovich and Sunyaev, 1969; Sunyaev and Zeldovich, 1970; Illarionov and Siunjaev, 1975; Sunyaev and Khatri, 2013). At redshifts below 1.5×10^4 , after thermalization decoupling, Compton and inverse Compton scattering produce the so-called " y distortions"⁴ (Zeldovich and Sunyaev, 1969; Sunyaev and Zeldovich, 1972). For $1.5 \times 10^4 < z < 2 \times 10^5$, we get a distortion spectrum that is intermediate between those of y and μ -type distortions (Khatri and Sunyaev, 2012).

The two different types of distortions have different frequency dependence and this can suggest the type (as well as redshift) of interactions that give rise to spectral distortions.

Assume PIXIE will have the sensitivity as described in Table 1

¹The branch of cosmology that makes detailed quantitative predictions and measurements of properties of the Universe

²You can read more at <https://ned.ipac.caltech.edu/level5/Sept05/Gawiser2/Gawiser1.html>

³The chemical potential μ is a quantity that expresses energy that is absorbed or released when the number of particles - in this case, photons - in a system changes. Here, it is measured in units of kT and is thus dimensionless.

⁴ y is a parameter that expresses by how much a photon will change its energy due to repeated scatterings in a medium of finite extent.

| | | | | |
|---------------------------|--------|--------|--------|--------|
| Frequency [GHz] | 14.4 | 374.6 | 734.9 | 1095.1 |
| PIXIE Sensitivity [Jy/sr] | 3.6035 | 3.7700 | 4.2859 | 5.1598 |

Table 1: PIXIE Sensitivity (Jy refers to the unit Jansky)

Spectral Distortions Modeling

y distortion PIXIE can shed light on the history of star formation by looking at the spectral distortions produced during the epoch of reionization. The CMB photons undergo inverse Compton scattering off the gas that has been ionized by early stars. These distortions are parametrized by y which expresses the mean number of scatterings times the average energy change that a photon suffers in a scattering. The respective intensity contribution is given by:

$$\Delta I_{\nu,y} = I_0 \frac{x^4 e^x}{(e^x - 1)^2} \left[x \coth\left(\frac{x}{2}\right) - 4 \right] y, \quad (1)$$

where $I_0 = \frac{2h}{c^2} \left(\frac{kT_0}{h}\right)^3 = 270 \text{ MJy/sr}$ for $T_0 = 2.726 \text{ K}$, and $x = \frac{h\nu}{kT_0}$. We take $y = 1.7 \times 10^{-6}$ for our fiducial value (assumed value for comparison).

μ distortion The amplitude of density fluctuations during inflation translates into energy injection in the CMB for redshifts $10^5 < z < 10^6$ which leads to μ distortions characterized by the chemical potential, with intensity given by

$$\Delta I_{\nu,\mu} = I_0 \frac{x^4 e^x}{(e^x - 1)^2} \left[\frac{1}{\beta} - \frac{1}{x} \right] \mu, \quad (2)$$

where $I_0 = \frac{2h}{c^2} \left(\frac{kT_0}{h}\right)^3 = 270 \text{ MJy/sr}$ for $T_0 = 2.726 \text{ K}$, $x = \frac{h\nu}{kT_0}$ and $\beta = 2.1923$.

A suitable fiducial value for μ is $\mu = 2 \times 10^{-8}$ (Abitbol et al., 2017).

Your task, as a team, is to calculate the expected standard deviations of the y and μ parameters, if PIXIE were to try to measure them in the bands given in Table 1. This will tell us if the y and μ distortions will be detectable, assuming the fiducial values given above.

In order to calculate that, you can use the matrix⁵:

$$F_{ij} = \sum_{a,b} \left(\frac{\partial(\Delta I_\nu)}{\partial p_i} \right)_a C_{ab}^{-1} \left(\frac{\partial(\Delta I_\nu)}{\partial p_j} \right)_b \quad (3)$$

called the Fisher Information Matrix (see for example Verde, 2010). a, b are indexing frequency, and p_i, p_j are parameters of the model (in our case, the two free parameters for the spectral distortions, y and μ , all other parameters assumed fixed). C_{ab}^{-1} is the inverse of the covariance matrix of the experiment. Inverting the F_{ij} matrix and taking the square root of the diagonal gives us the standard deviations expected for each parameter.

- (a) **(18 points)** Make a plot of $\Delta I_{\nu,y}$ and $\Delta I_{\nu,\mu}$ over the range of frequencies 14.4GHz to 1100.1 GHz, assuming the fiducial values for y and μ .

⁵For students not familiar with matrix notation and/or partial derivatives, a basic introduction for these can be found in any good introductory textbook on calculus / analysis

- (b) **(10 points)** Calculate the analytical forms of $\frac{\partial \Delta I_{\nu, \mu}}{\partial \mu}$ and $\frac{\partial \Delta I_{\nu, y}}{\partial y}$. Then, evaluate them at the 4 frequencies given in Table 1.
- (c) **(8 points)** Calculate the covariance matrix C_{ab} for PIXIE sensitivity in frequency bands using Table 1. The covariance matrix has the variance (square of the sensitivity) of the frequency bands on the diagonal, and the covariance of the bands on the other entries. In this problem, assume the frequency bands are uncorrelated and their covariance is thus 0.
- (d) **(4 points)** Calculate the inverse C_{ab}^{-1} of the PIXIE covariance matrix.
- (e) **(20 points)** Assuming PIXIE is trying to model only the y and μ distortions, and ignoring other foregrounds, **calculate** the Fisher Information Matrix, and the standard deviations for the parameters y and μ .
- (i) Would the y distortion be detectable with PIXIE?
- (ii) Would the μ distortion be detectable with PIXIE?

For the distortion to be detectable, the standard deviation on the parameter should be less than the value of the parameter.

Notes:

- Useful Constants and Units
 - $h = 6.626\,070\,04 \times 10^{-34} \text{ J s}$
 - $k = 1.380\,648\,52 \times 10^{-23} \text{ J K}^{-1}$
 - $1 \text{ Jy} = 10^{-26} \text{ W/m}^2/\text{Hz}$
- In this analysis, for each given frequency, the only free parameters of the functions $\Delta I_{\nu, y}$ and $\Delta I_{\nu, \mu}$ are y and μ respectively.
- A partial derivative $\frac{\partial f}{\partial x}$ of a multi-variable function $f(x, y, z, \dots)$ with respect to a single variable x is the process of calculating derivative with respect to the said variable x , while all the other variables are kept constant.
- In reality, PIXIE will have to model other foregrounds as well, such as the foregrounds coming from the synchrotron emission in the galaxy, the thermal radiation from the dust present in the interstellar medium, the free-free emission, cosmic infrared background etc. But we will ignore these contributions for the purpose of the problem.

References

- M. H. Abitbol, J. Chluba, J. C. Hill, and B. R. Johnson. Prospects for measuring cosmic microwave background spectral distortions in the presence of foregrounds. *Monthly Notices of the Royal Astronomical Society, Volume 471, Issue 1, p.1126-1140*, 471:1126–1140, oct 2017. doi: 10.1093/mnras/stx1653. URL <https://ui.adsabs.harvard.edu/abs/2017MNRAS.471.1126A>.
- J. Chluba. Science with CMB spectral distortions, may 2014. URL <https://ui.adsabs.harvard.edu/abs/2014arXiv1405.6938C>.
- J. Chluba and D. Jeong. Teasing bits of information out of the CMB energy spectrum. *Monthly Notices of the Royal Astronomical Society, Volume 438, Issue 3, p.2065-2082*, 438:2065–2082, mar 2014. doi: 10.1093/mnras/stt2327. URL <https://ui.adsabs.harvard.edu/abs/2014MNRAS.438.2065C>.

- D. J. Fixsen. The Temperature of the Cosmic Microwave Background. *The Astrophysical Journal*, Volume 707, Issue 2, pp. 916-920 (2009)., 707:916–920, dec 2009. doi: 10.1088/0004-637X/707/2/916. URL <https://ui.adsabs.harvard.edu/abs/2009ApJ...707..916F>.
- D. J. Fixsen, E. S. Cheng, J. M. Gales, J. C. Mather, R. A. Shafer, and E. L. Wright. The Cosmic Microwave Background Spectrum from the Full COBE FIRAS Data Set. *Astrophysical Journal* v.473, p.576, 473:576, dec 1996. doi: 10.1086/178173. URL <https://ui.adsabs.harvard.edu/abs/1996ApJ...473..576F>.
- J. C. Hill, N. Battaglia, J. Chluba, S. Ferraro, E. Schaan, and D. N. Spergel. Taking the Universe’s Temperature with Spectral Distortions of the Cosmic Microwave Background. *Physical Review Letters*, Volume 115, Issue 26, id.261301, 115:261301, dec 2015. doi: 10.1103/PhysRevLett.115.261301. URL <https://ui.adsabs.harvard.edu/abs/2015PhRvL.115z1301H>.
- A. F. Illarionov and R. A. Sunyaev. Comptonization, characteristic radiation spectra, and thermal balance of low-density plasma. *Soviet Astronomy*, vol. 18, Jan.-Feb. 1975, p. 413-419. Translation., 18:413–419, feb 1975. URL <https://ui.adsabs.harvard.edu/abs/1975SvA....18..413I>.
- R. Khatri and R. A. Sunyaev. Beyond γ and μ : the shape of the CMB spectral distortions in the intermediate epoch, $1.5 \tilde{\text{A}} \leq \nu \leq 105 \tilde{\text{A}}$. *Journal of Cosmology and Astroparticle Physics*, Issue 09, article id. 016 (2012)., 2012:16, sep 2012. doi: 10.1088/1475-7516/2012/09/016. URL <https://ui.adsabs.harvard.edu/abs/2012JCAP...09..016K>.
- A. Kogut, D. J. Fixsen, D. T. Chuss, J. Dotson, E. Dwek, M. Halpern, G. F. Hinshaw, S. M. Meyer, S. H. Moseley, M. D. Seiffert, D. N. Spergel, and E. J. Wollack. The Primordial Inflation Explorer (PIXIE): a nulling polarimeter for cosmic microwave background observations. *Journal of Cosmology and Astroparticle Physics*, Issue 07, id. 025 (2011)., 2011:25, jul 2011. doi: 10.1088/1475-7516/2011/07/025. URL <https://ui.adsabs.harvard.edu/abs/2011JCAP...07..025K>.
- A. Kogut, J. Chluba, D. J. Fixsen, S. Meyer, and D. Spergel. The Primordial Inflation Explorer (PIXIE). In *Proceedings of the SPIE*, Volume 9904, id. 99040W 23 pp. (2016)., volume 9904, page 9904W, jul 2016. doi: 10.1117/12.2231090. URL <https://ui.adsabs.harvard.edu/abs/2016SPIE.9904E..0WK>.
- J. C. Mather, E. S. Cheng, D. A. Cottingham, R. E. Eplee Jr., D. J. Fixsen, T. Hewagama, R. B. Isaacman, K. A. Jensen, S. S. Meyer, P. D. Noerdlinger, S. M. Read, L. P. Rosen, R. A. Shafer, E. L. Wright, C. L. Bennett, N. W. Boggess, M. G. Hauser, T. Kelsall, S. H. Moseley Jr., R. F. Silverberg, G. F. Smoot, R. Weiss, and D. T. Wilkinson. Measurement of the Cosmic Microwave Background Spectrum by the COBE FIRAS Instrument. *Astrophysical Journal* v.420, p.439, 420:439, jan 1994. ISSN 0004-637X. doi: 10.1086/173574. URL <https://ui.adsabs.harvard.edu/abs/1994ApJ...420..439M>.
- R. A. Sunyaev and R. Khatri. Unavoidable CMB Spectral Features and Blackbody Photosphere of Our Universe. *International Journal of Modern Physics D*, Volume 22, Issue 7, id. 1330014, 22:1330014, jun 2013. doi: 10.1142/S0218271813300140. URL <https://ui.adsabs.harvard.edu/abs/2013IJMPD...2230014S>.
- R. A. Sunyaev and Y. B. Zeldovich. The interaction of matter and radiation in the hot model of the Universe, II. *Astrophysics and Space Science*, 7:20–30, apr 1970. ISSN 0004-640X. doi: 10.1007/BF00653472. URL <https://ui.adsabs.harvard.edu/abs/1970ApJ...7...20S>.
- R. A. Sunyaev and Y. B. Zeldovich. Formation of Clusters of Galaxies; Protocluster Fragmentation and Intergalactic Gas Heating. *Astronomy and Astrophysics*, 20:189, aug 1972. ISSN 0004-6361. URL <https://ui.adsabs.harvard.edu/abs/1972A&A...20..189S>.

- H. Tashiro. CMB spectral distortions and energy release in the early universe. *Progress of Theoretical and Experimental Physics, Volume 2014, Issue 6, id.06B10727 pp.*, 2014:06B107, jun 2014. doi: 10.1093/ptep/ptu066. URL <https://ui.adsabs.harvard.edu/abs/2014PTEP.2014fB107T>.
- L. Verde. Statistical Methods in Cosmology. In *Lectures on Cosmology Accelerated Expansion of the Universe by Georg Wolschin, Lecture Notes in Physics vol. 800, Springer Berlin / Heidelberg, ISSN 1616-6361; ISBN 978-3-642-10597-5, pp.147-177*, volume 800, pages 147–177. mar 2010. ISBN 1616-6361. doi: 10.1007/978-3-642-10598-2_4. URL <https://ui.adsabs.harvard.edu/abs/2010LNP...800..147V>.
- Y. B. Zeldovich and R. A. Sunyaev. The Interaction of Matter and Radiation in a Hot-Model Universe. *Astrophysics and Space Science*, 4:301–316, jul 1969. ISSN 0004-640X. doi: 10.1007/BF00661821. URL <https://ui.adsabs.harvard.edu/abs/1969ApJSS...4..301Z>.